## Contents

Volume 110 1990

1	Adams, D. S., Keller, R. and Koehl, M. A. R. The mechanics of notochord elongation, straightening and stiffening in the embryo of <i>Xenopus laevis</i>	115
10	Yu, CK. C., Tsui, LC. and Breitman, M. L. Homologous and heterologous enhancers modulate spatial expression but not cell-type specificity of the murine $\gamma$ F-crystallin promoter	131
19	Payre, F., Noselli, S., Lefrère, V. and Vincent, A. The closely related $Drosophila\ sry\ \beta$ and $sry\ \delta$ zinc finger proteins show differential embryonic expression and distinct patterns of binding sites on polytene chromosomes	141
31	Lee, J. E., Pintar, J. and Efstratiadis, A. Pattern of the insulin-like growth factor II gene expression during early mouse embryogenesis	151
41	Macdonald, P. M.  bicoid mRNA localization signal: phylogenetic conservation of function and RNA secondary structure	161
51	Watson, M. A. and Milbrandt, J. Expression of the nerve growth factor-regulated NGFI-A and NGFI-B genes in the developing rat	173
	Martin-Morris, L. E. and White, K. The <i>Drosophila</i> transcript encoded by the $\beta$ -amyloid protein precursor-like gene is restricted to the nervous system	185
63	Sadaghiani, B. and Vielkind, J. R. Distribution and migration pathways of HNK-1-immunoreactive neural crest cells in teleost fish embryos	197
73	Adler, R. R., Brenner, C. A. and Werb, Z. Expression of extracellular matrix-degrading metalloproteinases and metalloproteinase inhibitors is developmentally regulated during endoderm differentiation of embryonal carcinoma	
97	Corps, A. N., Brigstock, D. R., Littlewood, C. J. and Brown, K. D. Receptors for epidermal growth factor and	211
105	trophoderm of the pig  Musci, T. S. and Mullen, R. J.  All-or-none craniorachischisis in Loop-tail mutant mouse chimeras	221
	19 31 41 51 63 85 97	The mechanics of notochord elongation, straightening and stiffening in the embryo of <i>Xenopus laevis</i> Yu, CK. C., Tsui, LC. and Breitman, M. L. Homologous and heterologous enhancers modulate spatial expression but not cell-type specificity of the murine γF-crystallin promoter  Payre, F., Noselli, S., Lefrère, V. and Vincent, A. The closely related <i>Drosophila sry β</i> and <i>sry δ</i> zinc finger proteins show differential embryonic expression and distinct patterns of binding sites on polytene chromosomes  Lee, J. E., Pintar, J. and Efstratiadis, A. Pattern of the insulin-like growth factor II gene expression during early mouse embryogenesis  Macdonald, P. M. bicoid mRNA localization signal: phylogenetic conservation of function and RNA secondary structure  Watson, M. A. and Milbrandt, J. Expression of the nerve growth factor-regulated NGFI-A and NGFI-B genes in the developing rat  Martin-Morris, L. E. and White, K. The <i>Drosophila</i> transcript encoded by the β-amyloid protein precursor-like gene is restricted to the nervous system  Sadaghiani, B. and Vielkind, J. R. Distribution and migration pathways of HNK-1-immunoreactive neural crest cells in teleost fish embryos  Adler, R. R., Brenner, C. A. and Werb, Z. Expression of extracellular matrix-degrading metalloproteinases and metalloproteinase inhibitors is developmentally regulated during endoderm differentiation of embryonal carcinoma cells  Corps, A. N., Brigstock, D. R., Littlewood, C. J. and Brown, K. D.  Receptors for epidermal growth factor and insulin-like growth factor-I on preimplantation trophoderm of the pig

Birr, C. A., Fristrom, D., King, D. S. and Fristrom, J. W. Ecdysone-dependent proteolysis of an apical surface glycoprotein may play a role in imaginal disc morphogenesis in <i>Drosophila</i>	239	Altmann, G. G. and Quaroni, A. Behavior of fetal intestinal organ culture explanted onto a collagen substratum  Vaessen, MJ., Meijers, J. H. C., Bootsma, D.	353
Hake, L. E., Alcivar, A. A. and Hecht, N. B. Changes in mRNA length accompany translational regulation of the somatic and testisspecific cytochrome c genes during		and van Kessel, A. G. The cellular retinoic-acid-binding protein is expressed in tissues associated with retinoic-acid- induced malformations	371
spermatogenesis in the mouse	249	Ohneda, O., Yanai, N. and Obinata, M. Microenvironment created by stromal cells is	
Tomlinson, C. R., Kozlowski, M. T. and Klein, W. H. Ectoderm nuclei from sea urchin embryos contain		essential for a rapid expansion of erythroid cells in mouse fetal liver	379
a Spec-DNA binding protein similar to the vertebrate transcription factor USF	259	Renfree, M. B., Robinson, E. S., Short, R. V. and Vandeberg, J. L.	
Takeda, H., Suematsu, N. and Mizuno, T. Transcription of prostatic steroid binding protein (PSBP) gene is induced by		Mammary glands in male marsupials: I. Primordia in neonatal opossums <i>Didelphis virginiana</i> and <i>Monodelphis domestica</i>	385
epithelial-mesenchymal interaction	273	Tesarik, J., Drahorad, J., Testart, J. and Mendoza, C.	
Hauxwell, A. J., Corke, F. M. K., Hedley, C. L. and Wang, T. L. Storage protein gene expression is localised to regions lacking mitotic activity in developing pea		Acrosin activation follows its surface exposure and precedes membrane fusion in human sperm acrosome reaction	391
embryos. An analysis of seed development in Pisum sativum XIV	283	Wehrle, B. and Chiquet, M. Tenascin is accumulated along developing	
Hidalgo, A. and Ingham, P. W. Cell patterning in the <i>Drosophila</i> segment: spatial regulation of the segment polarity gene <i>patched</i>	291	peripheral nerves and allows neurite outgrowth in vitro	401
Kwong, L., Xie, Y., Daniel, J., Robbins, S. M. and Weeks, G.  A Dictyostelium morphogen that is essential for stalk cell formation is generated by a subpopulation of prestalk cells	303	Charlebois, T. S., Henry, J. J. and Grainger, R. M. Differential cytokeratin gene expression reveals early dorsal-ventral regionalization in chick mesoderm	417
Sullivan, W., Minden, J. S. and Alberts, B. M. daughterless-abo-like, a Drosophila maternal-effect mutation that exhibits abnormal centrosome separation during the late blastoderm divisions	311	Ovsenek, N. and Heikkila, J. J. DNA sequence-specific binding activity of the heat-shock transcription factor is heat-inducible before the midblastula transition of early <i>Xenopus</i> development	427
Hemmati-Brivanlou, A., Frank, D., Bolce, M. E., Brown, R. D., Sive, H. L. and Harland, R. M. Localization of specific mRNAs in <i>Xenopus</i> embryos by whole-mount <i>in situ</i> hybridization	325	Albano, R. M., Godsave, S. F., Huylebroeck, D., van Nimmen, K., Isaacs, H. V., Slack, J. M. W. and Smith, J. C.  A mesoderm-inducing factor produced by	
Ichijo, H., Fujita, S., Matsuno, T. and Nakamura, H.	020	WEHI-3 murine myelomonocytic leukemia cells is activin A	435
Rotation of the tectal primordium reveals plasticity of target recognition in retinotectal projection	331	Gatherer, D., ten Dijke, P., Baird, D. T. and Akhurst, R. J. Expression of TGF- $\beta$ isoforms during first trimester human embryogenesis	445
Dencker, L., Annerwall, E., Busch, C. and Eriksson, U. Localization of specific retinoid-binding sites and		Otte, A. P., Kramer, I. M., Mannesse, M., Lambrechts, C. and Durston, A. J.	773
expression of cellular retinoic-acid-binding protein (CRABP) in the early mouse embryo	343	Characterization of protein kinase C in early Xenopus embryogenesis	461

Bennett, D. C., Huszar, D., Laipis, P. J., Jaenisch, R. and Jackson, I. J. Phenotypic rescue of mutant brown melanocytes by a retrovirus carrying a wild-type tyrosinase- related protein gene	471	Huang, J. Q., Turbide, C., Daniels, E., Jothy, S. and Beauchemin, N. Spatiotemporal expression of murine carcinoembryonic antigen (CEA) gene family members during mouse embryogenesis	573
Thompson, E. M., Fleming, K. A., Evans, D. J., Fundele, R., Surani, M. A. and Wright, N. A. Gastric endocrine cells share a clonal origin with other gut cell lineages	477	Frohman, M. A., Boyle, M. and Martin, G. R. Isolation of the mouse <i>Hox-2.9</i> gene; analysis of embryonic expression suggests that positional information along the anterior–posterior axis is	500
Kucera, J. and Walro, J. M. Treatment with $\beta$ bungarotoxin blocks muscle spindle formation in fetal rats	483	Pelton, R. W., Dickinson, M. E., Moses, H. L. and Hogan, B. L. M.	589
Metcalfe, W. K., Myers, P. Z., Trevarrow, B., Bass, M. B. and Kimmel, C. B. Primary neurons that express the L2/HNK-1		In situ hybridization analysis of TGF $\beta$ 3 RNA expression during mouse development: comparative studies with TGF $\beta$ 1 and $\beta$ 2	609
carbohydrate during early development in the zebrafish  Tuan, R. S. and Bigioni, N.	491	Casanova, J. Pattern formation under the control of the terminal system in the <i>Drosophila</i> embryo	621
Ca <sup>2+</sup> -activated ATPase of the mouse chorioallantoic placenta: developmental expression, characterization and cytohistochemical localization	505	Erselius, J. R., Goulding, M. D. and Gruss, P. Structure and expression pattern of the murine <i>Hox-3.2</i> gene	629
Harper, S. and Davies, A. M. NGF mRNA expression in developing cutaneous epithelium related to innervation density	515	Plachov, D., Chowdhury, K., Walther, C., Simon, D., Guenet, JL. and Gruss, P. <i>Pax8</i> , a murine paired box gene expressed in the developing excretory system and thyroid gland	643
Ginsburg, M., Snow, M. H. L. and McLaren, A. Primordial germ cells in the mouse embryo during gastrulation	521	ESSAY IN DEVELOPMENT Yamada, T.	045
Nadon, N. L., Duncan, I. D. and Hudson, L. D. A point mutation in the proteolipid protein gene of the 'shaking pup' interrupts oligodendrocyte development	529	Regulations in the induction of the organized neural system in amphibian embryos  CONTRIBUTED PAPERS Bodmer, R., Jan, L. Y. and Jan, Y. N.	653
Pullar, D., te Kronnie, G., Peiris, D., Taverne, N., Jeacock, M. K., Stroband, H. W. J. and Shperd, D. A. L.		A new homeobox-containing gene, msh-2, is transiently expressed early during mesoderm formation of <i>Drosophila</i>	661
Morphological and radiochemical evidence for the metabolism of exogenous proteins by the preimplantation sheep blastocyst	539	Ihara, S., Motobayashi, Y., Nagao, E. and Kistler, A. Ontogenetic transition of wound healing pattern	
Saggin, L., Gorza, L., Ausoni, S. and Schiaffino, S. Cardiac troponin T in developing, regenerating		in rat skin occurring at the fetal stage  Nardi, J. B.	671
and denervated rat skeletal muscle  Hahnel, A. C., Rappolee, D. A., Millan, J. L.,  Manes, T., Ziomek, C. A., Theodosiou, N. G.,	547	Expression of a surface epitope on cells that link branches in the tracheal network of <i>Manduca sexta</i>	681
Werb, Z., Pedersen, R. A. and Schultz, G. A. Two alkaline phosphatase genes are expressed during early development in the mouse embryo	555	Baetge, G., Schneider, K. A. and Gershon, M. D. Development and persistence of catecholaminergic neurons in cultured explants of fetal murine vagus nerves and bowel	689
Tanaka, H., Kinutani, M., Agata, A., Takashima, Y. and Obata, K. Pathfinding during spinal tract formation in the chick-quail chimera analysed by species-specific		Marin, M. L., Tobias, M. L. and Kelley, D. B. Hormone-sensitive stages in the sexual differentiation of laryngeal muscle fiber number	
monoclonal antibodies	565	in Xenopus laevis	703

Austin, C. P. and Cepko, C. L. Cellular migration patterns in the developing mouse cerebral cortex	713	Henry, J. J., Wray, G. A. and Raff, R. A.  The dorsoventral axis is specified prior to first cleavage in the direct developing sea urchin	075
Izpisùa-Belmonte, JC., Dollé, P., Renucci, A., Zappavigna, V., Falkenstein, H. and Duboule, D. Primary structure and embryonic expression pattern of the mouse <i>Hox-4.3</i> homeobox gene	733	Vize, P. D., Vaughan, A. and Krieg, P. Expression of the N-myc proto-oncogene during the early development of Xenopus laevis	875 885
Hayes, W. P. and Loh, Y. P. Correlated onset and patterning of proopiomelanocortin gene expression in embryonic <i>Xenopus</i> brain and pituitary	747	Flanders, D. J., Rawlins, D. J., Shaw, P. J. and Lloyd, C. W.  Re-establishment of the interphase microtubule array in vacuolated plant cells, studied by confocal microscopy and 3-D imaging	897
Warrior, R. and Levine, M.  Dose-dependent regulation of pair-rule stripes by gap proteins and the initiation of segment polarity	759	Haenlin, M., Kramatschek, B. and Campos-Ortega, J. A. The pattern of transcription of the neurogenic	
Smith, R. C., Bement, W. M., Dersch, M. A., Dworkin-Rastl, E., Dworkin, M. B.		gene Delta of Drosophila melanogaster	905
and Capco, D. G. Nonspecific effects of oligodeoxynucleotide injection in <i>Xenopus</i> oocytes: a reevaluation of previous D7 mRNA ablation experiments	769	Tear, G., Akam, M. and Martinez-Arias, A. Isolation of an <i>abdominal-A</i> gene from the locust <i>Schistocerca gregaria</i> and its expression during early embryogenesis	915
del Pino, E. M. and Loor-Vela, S. The pattern of early cleavage of the marsupial frog Gastrotheca riobambae	781	Simpson, P. and Carteret, C. Proneural clusters: equivalence groups in the epithelium of <i>Drosophila</i>	927
<b>Bate, M.</b> The embryonic development of larval muscles in <i>Drosophila</i>	791	Lackie, P. M., Zuber, C. and Roth, J. Polysialic acid and N-CAM localisation in embryonic rat kidney: mesenchymal and epithelial elements show different patterns of	
Brauer, P. R., Keller, K. M. and Keller, J. M. Concurrent reduction in the sulfation of heparan sulfate and basement membrane assembly in a cell model system	805	expression  Brown, N. A., McCarthy, A. and Wolpert, L. The development of handed asymmetry in	933
Nagy, A., Gócza, E., Diaz, E. M., Prideaux, V. R., Iványi, E., Markkula, M.		aggregation chimeras of situs inversus mutant and wild-type mouse embryo	949
and Rossant, J.  Embryonic stem cells alone are able to support fetal development in the mouse	815	Bassez, T., Paris, J., Omilli, F., Dorel, C. and Osborne, H. B. Post-transcriptional regulation of ornithine	
Klein, G., Ekblom, M., Fecker, L., Timpl, R. and Ekblom, P. Differential expression of laminin A and B chains		decarboxylase in <i>Xenopus laevis</i> oocytes  Harvey, M. B. and Kaye, P. L.  Insulin increases the cell number of the inner cell	955
during development of embryonic mouse organs	823	mass and stimulates morphological development of mouse blastocysts <i>in vitro</i>	963
Charpentier, G. and Magre, S. Masculinizing effect of testes on developing rat ovaries in organ culture	839	Shao, C. and Takagi, N. An extra maternally derived X chromosome is deleterious to early mouse development	969
Simpson, L. and Wieschaus, E. Zygotic activity of the <i>nullo</i> locus is required to stabilize the actin–myosin network during cellularization in <i>Drosophila</i>	851	Berks, M. and Kay, R. R. Combinatorial control of cell differentiation by cAMP and DIF-1 during development of Dictyostelium discoideum	977
Yost, H. J. Inhibition of proteoglycan synthesis eliminates left-right asymmetry in <i>Xenopus laevis</i> cardiac looping	865	Sylvester, A. W., Cande, W. Z. and Freeling, M. Division and differentiation during normal and liguleless-1 maize leaf development	985

ESSAY IN DEVELOPMENT		Kaufman, M. H. and Webb, S.	
Potten, C. S. and Loeffler, M. Stem cells: attributes, cycles, spirals, pitfalls and		Postimplantation development of tetraploid mouse embryos produced by electrofusion	1121
	1001	mouse emoryos produced by electrorusion	1121
		Dollé, P., Ruberte, E., Leroy, P., Morriss-Kay, G.	•
CONTRIBUTED PAPERS		and Chambon, P.	
Chuong, CM., Oliver, G., Ting, S. A., Jegalian, B. G., Chen, H. M. and De		Retinoic acid receptors and cellular retinoid	
Robertis, E. M.		binding proteins. I. A systematic study of their differential pattern of transcription during mouse	
Gradients of homeoproteins in developing feather		organogenesis	1133
buds	1021		2200
Pouton D. Dougouiban C. F. F. Hoffman F. M.		Pohl, T. M., Mattei, MG. and Rüther, U.	
Reuter, R., Panganiban, G. E. F., Hoffman, F. M. and Scott, M. P.	•	Evidence for allelism of the recessive insertional	
Homeotic genes regulate the spatial expression of		mutation <i>add</i> and the dominant mouse mutation <i>extra-toes</i> ( <i>Xt</i> )	1153
putative growth factors in the visceral mesoderm		extra-toes (Xt)	1133
of <i>Drosophila</i> embryos	1031	Vogels, R., de Graaf, W. and Deschamps, J.	
Demonstra C F F Deuten D Coatt M D		Expression of the murine homeobox-containing	
Panganiban, G. E. F., Reuter, R., Scott, M. P. and Hoffman, F. M.		gene Hox-2.3 suggests multiple time-dependent	44.50
A Drosophila growth factor homolog,		and tissue-specific roles during development	1159
decapentaplegic, regulates homeotic gene		Katzen, A. L., Kornberg, T. and Bishop, J. M.	
expression within and across germ layers during		Diverse expression of <i>dsrc29A</i> , a gene related to	
midgut morphogenesis	1041	src, during the life cycle of Drosophila	
Yuge, M., Kobayakawa, Y., Fujisue, M.		melanogaster	1169
and Yamana, K.		Town N. D. Horman, I. and William C. C.	
A cytoplasmic determinant for dorsal axis		Torpey, N. P., Heasman, J. and Wylie, C. C. Identification of vimentin and novel vimentin-	
formation in an early embryo of Xenopus laevis	1051	related proteins in <i>Xenopus</i> oocytes and early	
Manaya K Nagka K Pasmar D		embryos	1185
Manova, K., Nocka, K., Besmer, P. and Bachvarova, R. F.			
Gonadal expression of c-kit encoded at the W		Macias, A., Casanova, J. and Morata, G.	
locus of the mouse	1057	Expression and regulation of the <i>abd-A</i> gene of <i>Drosophila</i>	1197
N		Drosophina	1171
Nübler-Jung, K. and Mardini, B.  Insect epidermis: polarity patterns after grafting		Tilney, L. G., Cooke, T. J., Connelly, P. S.	
result from divergent cell adhesions between host		and Tilney, M. S.	
and graft tissue	1071	The distribution of plasmodesmata and its	
		relationship to morphogenesis in fern gametophytes	1209
Represa, J., Sanchez, A., Miner, C., Lewis, J.		gametophytes	1207
and Giraldez, F. Retinoic acid modulation of the early		Howard, K. R. and Struhl, G.	
development of the inner ear is associated with		Decoding positional information: regulation of	
the control of c-fos expression	1081	the pair-rule gene hairy	1223
Calana I Office W C Nation D D		Mackie, E. J., Trechsel, U. and Bruns, C.	
Schuger, L., O'Shea, K. S., Nelson, B. B.		Somatostatin receptors are restricted to a	
and Varani, J.  Organotypic arrangement of mouse embryonic		subpopulation of osteoblast-like cells during	
lung cells on a basement membrane extract:		endochondral bone formation	1233
involvement of laminin	1091	Tallamana Wanad DaAlaa D	
Thomas W A Cabactar A W		Lallemand, Y. and Brûlet, P.  An in situ assessment of the routes and extents of	f
Thomas, W. A., Schaefer, A. W. and Treadway, R. M. Jr		colonisation of the mouse embryo by embryonic	
Galactosyl transferase-dependence of neurite		stem cells and their descendants	1241
outgrowth on substratum-bound laminin	1101	D 00 I III III 40 II W 0 II	
Walaki I V and Dames I D A		Raff, J. W., Whitfield, W. G. F.	
Knight, J. K. and Raymond, P. A.  Time course of opsin expression in developing		and Glover, D. M. Two distinct mechanisms localise cyclin B	
rod photoreceptors	1115	transcripts in syncytial <i>Drosophila</i> embryos	1249

## Contents

Biggiogera, M., Bürki, K., Kaufmann, S. H., Shaper, J. H., Gas, N., Amalric, F. and Fakan, S Nucleolar distribution of proteins B23 and nucleolin in mouse preimplantation embryos as visualized by immunoelectron microscopy	1263	Gould, A. P., Lai, R. Y. K., Green, M. J. and White, R. A. H.  Blocking cell division does not remove the requirement for <i>Polycomb</i> function in <i>Drosophila</i> embryogenesis	1319
Stern, C. D., Ireland, G. W., Herrick, S. E., Gherardi, E., Gray, J., Perryman, M. and Stoker, M. Epithelial scatter factor and development of the chick embryonic axis	1271	Hortsch, M., Patel, N. H., Bieber, A. J., Traquina, Z. R. and Goodman, C. S. Drosophila neurotactin, a surface glycoprotein with homology to serine esterases, is dynamically expressed during embryogenesis	132
Kücherer-Ehret, A., Pottgiesser, J., Kreutzberg, G. W., Thoenen, H. and Edgar, D. Developmental loss of laminin from the interstitial extracellular matrix correlates with decreased laminin gene expression	1285	Nichols, J., Evans, E. P. and Smith, A. G. Establishment of germ-line-competent embryonic stem (ES) cells using differentiation inhibiting activity	134
Swann, K.  A cytosolic sperm factor stimulates repetitive calcium increases and mimics fertilization in hamster eggs	1295	Honda, H., Tanemura, M. and Yoshida, A. Estimation of neuroblast numbers in insect neurogenesis using the lateral inhibition hypothesis of cell differentiation	134
Schauer, I. E. and Wood, W. B. Early <i>C. elegans</i> embryos are transcriptionally		Index of Authors and Titles	135
active	1303	Subject Index	135

